

# CHEMOTHERAPY ORDERS – Docetaxel Weekly Schedule – Adult Solid Tumor

Date chemo. to be initiated: \_\_\_\_\_

Cycle #: \_\_\_\_\_

Weight: \_\_\_\_\_  lbs.  kg  
 actual  ideal

Height: \_\_\_\_\_  inches  cm  
BSA used for calculations: \_\_\_\_\_ m2

**LABS: CBC, DIFF, PLATELETS, CMP or BMP (circle appropriate one).**

For cycle 2 and subsequent cycles, are doses identical to previous cycle?  Yes  No

If no, indicate reason for adjustment: \_\_\_\_\_.

Dose reduction: \_\_\_\_\_ %      Dose increase: \_\_\_\_\_ %

**Treat only if:** GRANULOCYTES greater than or equal to 1000  
PLATELETS greater than or equal to 100,000  
WBC greater than or equal to 3000

**Premedicate** 30 minutes prior to chemotherapy with palonosetron 0.25 mg IV and dexamethasone 20 mg po.

**Docetaxel (Taxotere)** \_\_\_\_\_ mg/m2 (usual dose 30-40 mg/m2) x \_\_\_\_\_ m2 = \_\_\_\_\_ mg in 250 ml D5W IV over 60 minutes.

If counts are adequate, treat weekly x \_\_\_\_\_ weeks with a \_\_\_\_\_ week rest. (Note that the usual regimen is either 3 weeks with a 1 week rest or 6 weeks with a 2 week rest.) Treat today, and repeat on \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_.

Monitor vital signs at beginning of infusion and at the end of the infusion.

**Reason for any dose adjustment:**

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

MD

Signature: \_\_\_\_\_

Printed

Name: \_\_\_\_\_

Pager #: \_\_\_\_\_



MR.ORDERS

SVPOD-036 (6/07)



ST. VINCENT CHARITY  
MEDICAL CENTER

2351 EAST 22ND STREET  
CLEVELAND, OH 44115  
stvincentcharity.com

A Ministry of the Sisters of Charity Health System

PATIENT LABEL