

CHEMOTHERAPY ORDERS - Docetaxel (Every 21 Days) - ADULT SOLID TUMOR

Date chemo to be initiated: _____ Cycle #: _____

Weight: _____ lbs. Kg Height: _____ inches cm
_____ actual ideal **BSA used for calculations:** _____ m2

LABS: CBC, DIFF, PLATELETS, CMP.

For cycle 2 and subsequent cycles, are doses identical to previous cycle? Yes No

If no, indicate reason for adjustment: _____

Dose reduction: _____% Dose increase: _____%

Treat only if: WBC greater than or equal to 3000
Granulocytes greater than or equal to 1500 Platelets greater than or equal to 100,000

PREMEDICATE 30 minutes prior to chemotherapy with palonosetron 0.25 mg IV, dexamethasone 20 mg po.

Docetaxel (Taxotere) _____ mg/m2 (usual dose 75-100 mg/m2) x _____ m2 = _____ mg in 250 mL D5W IV over 60 minutes. If total dose is greater than 240 mg, then dilute in 500 mL of D5W. (Note that for second line chemotherapy, the usual starting dose is 75 mg/m2).

Dexamethasone 4 mg po BID x 4 days starting from day 1 each cycle.
(Patient needs prescription for this to go home with).

Monitor vital signs at beginning of infusion and at the end of infusion.

REASON FOR DOSE ADJUSTMENT:

MD Signature: _____ Printed Name: _____ Pager #: _____

Patient Name: _____



MR.ORDERS

SVPOD-038 (6/07)



ST. VINCENT CHARITY
MEDICAL CENTER

2351 EAST 22ND STREET
CLEVELAND, OH 44115
stvincentcharity.com

A Ministry of the Sisters of Charity Health System

PATIENT LABEL