

MANAGEMENT OF SEPTIC SHOCK: EARLY GOAL DIRECTED THERAPY MUST START WITHIN 6 HOURS

Central venous access (triple lumen catheter)

inserted by: Surgical House Officer Resident
 Chest X ray to confirm placement of TLC (not for femoral placement)

Intubation: Yes No
 Chest X ray to confirm placement of ET tube

Initial ventilator settings

ABG 30 minutes after intubation

Modes: A/C Synchronized Intermittent Mandatory ventilation (SIMV)

Vt (tidal volume) _____ RR (respiratory rate) _____

PEEP _____ (cmH₂O) No PEEP

FiO₂ _____ %

SVO₂ continuous cardiac output monitor: Yes No

Arterial Line: Radial Brachial Femoral

inserted by: Surgical House Officer Resident

Resuscitation goals:

- Mean arterial pressure MAP greater than or equal to 65
- CVP greater than or equal to 8
- Urine output greater than or equal to 30 mL/hr
- Central venous oxygen saturation ScvO₂ greater than or equal to 70%. If venous oxygen saturation target is not achieved call M.D.

Consider further fluid IV bolus _____ mL of NS or 1/2 NS over _____ .

Type and cross match _____ Units of PRBC

Transfuse PRBC to keep hematocrit greater than or equal to 30%

Physician Printed Name: _____ Date: _____ Time: _____

Physician's Signature: _____

PROHIBITED ABBREVIATION	REQUIRED TERM	PROHIBITED ABBREVIATION	REQUIRED TERM
ug	Write Microgram	1.0	Write 1. Do not use zero after decimal point
qd, q.d.	Write Daily	Zero after decimal point	
qod	Write Every Other Day or Every 48 hrs	MS	Write Morphine
U or u	Write Units	MgSO ₄ , MSO ₄	Write Magnesium sulfate or Morphine sulfate
.5	Write 0.5 - make sure you use preceding 0	IU	Write International units
No zero before decimal point		OS, OD, OU	Write Left or right eye or both eyes
		AS, AD, AU	Write Left or right ear or both ears



ST. VINCENT CHARITY
MEDICAL CENTER

2351 EAST 22ND STREET
CLEVELAND, OH 44115
stvincentcharity.com

A Ministry of the Sisters of Charity Health System

PATIENT LABEL

MANAGEMENT OF SEPTIC SHOCK: EARLY GOAL DIRECTED THERAPY MUST START WITHIN 6 HOURS

Source identification of infection:

- Blood cultures and sensitivity x 2 to be drawn 15 minutes apart Urine analysis
- Urine culture & sensitivity Sputum culture and sensitivity
- Wound culture and sensitivity: Site of wound _____
Site of wound _____
- Other cultures and sensitivities: _____ (site)
_____ (site)

Antibiotic therapy BEGIN AS SOON AS POSSIBLE AND WITHIN ONE HOUR

Attempt to draw blood cultures prior to antibiotic administration

1. Name _____ Dose _____ IV every _____ hour
2. Name _____ Dose _____ IV every _____ hour
3. Name _____ Dose _____ IV every _____ hour

Vasopressor: After appropriate fluid resuscitation

standard hospital concentrations

Titrate to maintain MAP greater than or equal to 65

- Norepinephrine (Levophed)
- Phenylephrine (Neosynephrine)
- Dopamine

Steroids (Consider for adult septic shock when hypotension responds poorly to adequate fluid resuscitation and vasopressors)

- Hydrocortisone (Solu-cortef) 50 mg IV every 6 hours
- Fludrocortisone (Florinef) 50 mcg SL daily

DVT Prophylaxis:

- Pneumatic compression stockings OR Heparin 5000 units sq every 8 hours

Stress Ulcer Prophylaxis:

- Famotidine (Pepcid) 20 mg IV every 12 hours

Physician Printed Name: _____ Date: _____ Time: _____

Physician's Signature: _____

PROHIBITED ABBREVIATION	REQUIRED TERM	PROHIBITED ABBREVIATION	REQUIRED TERM
ug	Write Microgram	1.0	Write 1. Do not use zero after decimal point
qd, q.d.	Write Daily	Zero after decimal point	
qod	Write Every Other Day or Every 48 hrs	MS	Write Morphine
U or u	Write Units	MgSO ₄ , MSO ₄	Write Magnesium sulfate or Morphine sulfate
.5	Write 0.5 - make sure you use preceding 0	IU	Write International units
No zero before decimal point		OS, OD, OU AS, AD, AU	Write Left or right eye or both eyes Write Left or right ear or both ears



ST. VINCENT CHARITY
MEDICAL CENTER

2351 EAST 22ND STREET
CLEVELAND, OH 44115
stvincentcharity.com

A Ministry of the Sisters of Charity Health System

PATIENT LABEL