

ORTHO/PODIATRY - PHYSICIAN ORDER - ADULT ANTIBIOTIC FOR SURGICAL PROPHYLAXIS

Date of Procedure _____	Drug Allergies _____
-------------------------	----------------------

Height _____ Weight _____

Review patient allergies prior to prescribing/administering medications. Please select appropriate boxes.

Procedure	Drug of choice:	* If Penicillin Allergy:
Orthopedic/ Podiatry	Preop <input type="checkbox"/> Cefazolin 1 Gm IV (if wt less than 80 kg) <input type="checkbox"/> Cefazolin 2 Gm IV (if wt greater than or equal to 80 kg)	Preop <input type="checkbox"/> Vancomycin 1 Gm IV
	<input type="checkbox"/> Vancomycin 1 Gm IV for patient at high risk for MRSA (Reason must be specified _____)* Check box in NOTE.	NOTE: High Risk for MRSA <input type="checkbox"/> Nursing home patient <input type="checkbox"/> Dialysis patient <input type="checkbox"/> Recent Hospitalization <input type="checkbox"/> Recurrent antibiotic treatment <input type="checkbox"/> Chronic Wound Care <input type="checkbox"/> Known Prior colonization Other _____
	Intraop <input type="checkbox"/> Repeat Cefazolin dose if surgery exceeds 3 hours	
	Post-Op <input type="checkbox"/> Cefazolin 1 Gm IV every 8 hours x 2 doses (if wt less than 80 kg) <input type="checkbox"/> Cefazolin 2 Gm IV every 8 hours x 2 doses (if wt greater than or equal to 80 kg)	
<input type="checkbox"/> Vancomycin 1 Gm IV every 12 hours x 1 dose for patient at high risk for MRSA		
Head and Neck	Preop <input type="checkbox"/> Cefazolin 1 Gm IV (if wt less than 80 kg) <input type="checkbox"/> Cefazolin 2 Gm IV (if wt greater than or equal to 80 kg)	Preop <input type="checkbox"/> Clindamycin 600 mg IV <u>AND</u> Gentamicin 1.5 mg/kg
	Post-Op <input type="checkbox"/> Cefazolin 1 Gm IV (if wt less than 80 kg) every 8 hours x 2 doses	Post-Op <input type="checkbox"/> Clindamycin 600 mg IV every 8 hours x 2 doses <u>AND</u> Gentamicin IV Pharmacy to dose
	<input type="checkbox"/> Cefazolin 2 Gm IV (if wt greater than or equal to 80 kg) every 8 hours x 2 doses	

Doses are adjusted according to renal function. Duration of prophylaxis is NOT to exceed 24 hours from surgery close time. **Overwritten orders will not be dispensed by pharmacy since they do not meet evidenced based guidelines unless suspected infection is documented here:** _____

Physician Signature: _____ Pager: _____ Date: _____ Time: _____



ST. VINCENT CHARITY
MEDICAL CENTER

2351 EAST 22ND STREET
CLEVELAND, OH 44115
stvincentcharity.com

A Ministry of the Sisters of Charity Health System



MR.ORDER

SVPOD-199 (9/09)

PATIENT LABEL